

## Comparative Effects Of Rabbits Dung, Npk 15:15:15 And Cow Dung On The Growth And Yield Of Pepper

Aderemi, F. T.

Adewoye, A. A

Horticulture and Landscape Technology Department,  
Federal College of Forestry, Ibadan, Nigeria

Aderemi A. M

Shaib-Rahim, H. O.

Roberts, A. E.

Agricultural Technology Department, Federal College of  
Forestry, Ibadan, Nigeria

**Abstract:** Field experiment was conducted at the Teaching and Research Farm of the department of Agriculture Technology, Federal College of Forestry, Jericho, Ibadan, Oyo state, to study the effects of rabbit dung, NPK 15:15:15 and cow dung on growth and yield of sweet pepper (*Capsicum annuum*). The experimental design was completely randomized design (CRD) with 4 treatments and 4 replicates. The following variables were measured for the period of 9 weeks: plant height, leaves production, stem girth, number of fruits. The result obtained from the average mean shows that T1 (rabbit dung- 17g) had the best performance in the plant height, leave production, stem girth, number of fruit and fruit weight (19.48, 19.00, 1.85, 3.00, and 24.35 respectively), while top soil has least performance (14.7, 14.25, 1.28, 1.00, 2.50 respectively). ANOVA table revealed that there are significant differences ( $ON < p < .05$ ) in plant height, stem diameter, and number of fruit at 5% level of probability, while there is no significant difference ( $p > = 0.05$ ) in leave production, at 5% level of probability. Based on the result, it is therefore recommended that rabbit dung T1 (17g) should be adopted since it support the increase in plant height, leave production, stem girth, number of fruit, weight of the fruit of sweet pepper (*Capsicum annuum*).

### I. INTRODUCTION

Sweet pepper (*Capsicum annuum*) is a species of the plant genus *Capsicum*, native to Southern North American and Northern South. The species is the most common and extensive cultivated of the five domesticated *Capsicum*. (Poulas, 1994). Sweet pepper is called 'popurika' in Japan, 'peperoni' in Switzerland, 'poivon' in France, 'piment' in Korea. (Azhar *et al.*, 2004). Ideal growing conditions for sweet pepper include warm soil, ideally 21- 29°C (70-84°F) that is kept moist but not waterlogged (Labeled *et al.*, 2013). Pepper is an important crop not only because of its economic importance but also for nutritional value of its fruit being a major source of natural colour and antioxidant compound (Howard *et al.*, 1994). The intake of this compound in food is an important health protecting factors; they have been recognized as being beneficial for prevention of widespread human disease, like cancer when taken in adequate amount. *Capsicum annuum* fruit are consumed in fresh, dried or

processed form. Non-pungent fruit, usually called sweet pepper are eaten in raw salads but more commonly cooked, fried or processed together with other foods. They are consumed in such quality per serving that they constitute a real table vegetable contributing to the nutritional value of the meal, also rich in antioxidants and vitamin C. Sweet pepper was discovered to be a good source of medicinal preparation for black vomit, gout and paralysis (Khan *et al.*, 2010). It was reported by Juroszek and Tsai (2009) that sweet pepper fruits are good sources of many essential nutrients, such as vitamins A, C, and E, carotenoids, minerals (e.g., calcium and iron), and other secondary plant compounds.

Although pepper can be grown all over Nigeria, the northern region between latitude 10° N and 12° N is the major area for production where an estimated 77,000 hectares of land under pepper cultivation yields about 695 000 metric tons (FAO, 2000). The lower yield of pepper can be attributed to low and inherent soil fertility and management practices, weeds infestation and diseases problems (Grubben and El-

Tahir, 2004). One of the panacea for improved production output of tropical crop production is use of fertilizers and organic matter maintenance (Agboola, 1982). Fertilization studies on pepper production is pertinent in southern part of Nigeria due to high rainfall intensity which always resulted in leaching, impoverishing the soil. Research also shows that organic manure application is more beneficial over the use of chemical fertilization in tropical crop production, sustainability and soil fertility management (Ansa and Woke, 2018). There are different views as to which performs better on pepper (between organic and inorganic fertilizer). Thus, the researchers embarked on this study to investigate the effects of Rabbits Dung, NPK 15:15:15 and Cow Dung on the Growth and Yield of Pepper.

## II. MATERIALS AND METHOD

### STUDY SITE

The experiment was conducted at the Teaching and Research Farm of the department of Agriculture Technology, Federal College of Forestry, Jericho, Ibadan, Oyo state. The college area lies between latitude 7°26'N and longitude 3°36'E. The climate conditions of the area is tropically dominated by rainfall pattern, ranging from 1400-1500mm with average temperature of about 31.2°C. The area experience two distinct season; dry season, usually from November — March and raining season, from April-October (FRIN, 2011).

### NURSERY PRACTICE

Enclosed 1 m by 1 m nursery beds of dark top soil were made. The beds were watered before the seeds were broadcasted evenly and watered. After emergence, some seedlings were removed so that strong healthy pepper seedling can be obtained for transplanting. Seedlings were watered daily and ready for transplant after 40 days.

### EXPERIMENTAL PROCEDURES

The manures (rabbit dung and cow dung) and top soil were taken to the laboratory for analysis after they have been dried under intense sunlight for two weeks and later grind into powdery form. The manure were measured and weighed using sensitive weighing balance and thereafter put inside a polythene bag. The manures were incorporated into the soil in the polythene pot and watered to allow commencement of mineralization and germination four (4) days before transplanting. The study adopted a 4 x 4 factorial experiment arranged in a randomized complete block design (RCBD) and replicated four times making it a total of 16 test samples. The treatments were (1) T1-17g of Rabbit dung/3.5kg of top soil (2) T2- 1.7g of N.P.K 15:15:15/3.5kg of top soil (3) T3-17g of cow dung/3.5kg of top soil (4) T4- 3.5kg of Top Soil serving as the control.

## DATA COLLECTION AND ANALYSIS

Data collection commenced two weeks after planting and continued on weekly basis till the end of the experiment. The parameters assessed includes; plant height with the aid of meter rule, number of leaves which was counted with the aid of hand weekly, numbers of fruit was counted per plant, Stem girth was measured with the aid of vernier caliper. All data collected were subjected to analysis of variance (ANOVA) and significant means was separated using Duncan multiple range test (DMRT).

## III. RESULT AND DISCUSSIONS

### EFFECT OF THE TREATMENTS ON PLANT HEIGHT

Table 1 shows the result of effect of the treatments on plant height. From the table, the results of the height of *Capsicum annuum* varied on weekly basis, the mean value at the end of the eight weeks of the experiments were reported with plants treated with treatment 1 (T1) having the highest plant height with mean value of (19.48 cm), followed closely by treatment 2 (T2) with mean value of (19.23 cm), and treatment (T3) with mean value of (18.45 cm). While plant treated with top soil (control) has the least performance with mean value of (14.7 cm).

Treatment	1	2	3	4	5	6	7	8	9	Average
T1	16.1 3a	17.0 0a	17.20 a	17.38	19.3 3b	20. 90a	21. 45a	21. 57a	24. 50a	19.4 8a
T2	14.1 3b	15.0 0b	15.43 ab	15.58 ab	22.5 5	22. 13a	21. 40a	25. 00a	23. 73a	19.2 3a
T3	13.6 5b	14.1 3b	14.53 b	14.78 b	20.9 5	20. 50a	22. 10a	23. 18a	22. 33a	18.4 5a
T4	13.7 b	14.0 0b	14.13 b	14.38 b	14.8 8	15. 03b	15. 23b	15. 50a	15. 53b	14.7 b
% CV	17.2 3	17.0 9	16.95	16.61	10.2 1	13. 57	12. 27	13. 53	15. 20	10.8 2

Table 1

### EFFECT OF THE TREATMENT ON THE STEM DIAMETER

The result presented in Table 2 shows that rabbit dung had the best performance in collar diameter with an average mean of 1.85, N.P.K 15:15:15 performed better in collar diameter with an average mean of 1.65 while treatment 4 (Control) had the least performance with an average mean of 1.28.

Treatment	1	2	3	4	5	6	7	8	9	Average
T1	0.98b	1.28a	1.43a	1.58a	1.88a	1.93a	2.38a	2.45a	2.68a	1.85a
T2	0.90b	1.10ab	1.25a	1.43a	1.55ab	1.83a	2.05ab	2.28a	2.55a	1.65a
T3	2.90a	1.00b	1.20a	1.35a	1.40b	1.68ab	1.60bc	1.45b	1.38b	1.35b
T4	0.93b	1.05ab	1.30a	1.45a	1.23b	1.35b	1.38c	1.68b	1.75b	1.28b
% CV	143.09	14.7	11.75	14.56	17.75	14.59	18.50	18.58	18.32	10.88

Table 2

### AVERAGE NUMBER AND WEIGHT OF FRUITS

Table 3 shows the effects of rabbit dung, NPK 15:15:15, cow dung and top soil on number and weight of fruit of *Capsicum annuum*. The result shows that fruit weight of the *Capsicum annuum* planted on soil treated with rabbit dung (T1) has the highest mean value of (24.35), followed by (T2)

NPK 15:15:15 with mean value of (9.93), and cow dung (T2) with mean value of (3.37), while fruit weight of the *Capsicum annuum* on the control soil has least performance with mean value of (2.50). The result of the number of fruits is not relatively difference except for treatment one that has highest number of fruits (3). The number and length of fruits increased with the application of rabbit dung which was significantly different among the treatments applied. The results could be as a result of higher number of leaves, flowers and fruiting buds which may have increased fruit production (Nweke *et al.*, 2014).

Treatment	No. of Fruits	Weight of Fruits (g)
T1	3.00a	24.35
T2	1.50b	9.93
T3	1.00b	3.37
T4	1.00b	2.50
%CV	30.77	42.62

Table 3: Average number and weight of fruits

#### IV. CONCLUSION

The result of the present study indicated that the application of rabbit manure had more effect on the growth and yield components of *Capsicum annuum* compared to NPK 15:15:15 and cow dung as the highest values were recorded in rabbit manure treated plots in all the parameters measured. Rabbit manure released enough nutrients which resulted in significant increase in growth and yield of *Capsicum annuum*; also it serves as a good source of soil amendment, and improvement of soil properties which in turn resulted in improved growth and yield.

#### REFERENCES

[1] Agboola, A.A., (1982). Organic manuring and green manuring in Tropical Agricultural systems. Proc. of 12

Inter. Cong. of Soil Science, New Delhi, India 8-16 February, pp: 198

- [2] Ansa J.E.O. and Woke C. (2018). Effect of Spacing and Poultry Manure Rates on Growth, Yield and Quality of Cayenne Pepper (*Capsicum frutescens*. L) in Southern Rain Forest of Nigeria. International Journal of Environment, Agriculture and Biotechnology (IJEAB), Vol-3, Issue-4, pp 1234- 1240.
- [3] Azhar A., Farooqi and B.S Screeramu, (2004). University press, p bk, Revised 344p ISBN: 9788173715044.
- [4] FAO. (2000). Fertilizers and their Use. 4th Edn., Food and Agriculture Organization, Rome, Pages: 34.
- [5] Grubben, G.J.H., & El-Tahir, I.M. (2004). *Capsicum annum* L. In: (Grubben, G.J.H. and Denton, O.A, Editors). Plant Resources of Tropical Africa 2. Vegetables. (pp.154-163). PROTA Foundation, Wageningen, Netherlands/CTA Wageningen, Netherlands.
- [6] Juroszek, P., & Tsai, H. H. (2009). Yields of organically grown sweet pepper cultivars and lines during the hot-wet and cool-dry season in the tropics. Horticultural Technology, 19(2), 418-422.
- [7] Khan, M. S. I., Roy, S. S., & Pall, K. K. (2010). Nitrogen and phosphorus efficiency on the growth and yield attributes of *Capsicum*. Academic Journal of Plant Sciences, 3(2), 71-78.
- [8] Labeled, V., Obeid H. and Ressay, K. (2013). Effect of Relative Humidity and Temperature on pepper. C Species
- [9] Nweke IA, Okoli PSO, Enyioko CO (2014) Effect of different rates of poultry droppings and plant spacing on soil chemical properties and yield of cucumber. Elixir Agriculture 70: 23934 – 23940.
- [10] Poulos JM (1994) Pepper breeding (*Capsicum* spp.): achievements, challenges and possibilities. PlantBreed Abstr 64:143–155